WINREVAIR™
(sotatercept-csrk) for injection 45 mg, 60 mg

WHAT IS *WINREVAIR*?

WINREVAIR is a prescription medicine for adults with pulmonary arterial hypertension (PAH). PAH is a type of high blood pressure in the arteries of your lungs.

WINREVAIR can improve your ability to exercise and perform normal activities with fewer symptoms, and reduce the risk of your physical condition and symptoms worsening, including lowering your risk of hospitalization for PAH, lung transplant, and death.

STARTING THE CONVERSATION ABOUT *WINREVAIR* WITH YOUR DOCTOR

This guide can help you talk with your doctor about the possibilities for you with WINREVAIR. Bring this with you to your next visit.

TIPS FOR TALKING WITH YOUR DOCTOR

- **Read the questions below, choose the ones you want to ask, and add your own.**
- **Be honest and open about how you're feeling and your treatment goals.**
- **If there's something you don't understand, ask your doctor or a member of your care team to explain it in a different way.**
- **If you have a care partner, consider bringing them with you. They can take notes and/or speak on your behalf.**

ABOUT *WINREVAIR* AND HOW IT WORKS

- What is WINREVAIR?
- Have you prescribed WINREVAIR to other patients with PAH? Is it an option for me?
- How does WINREVAIR work?

Notes:

UNDERSTANDING POTENTIAL BENEFITS AND RISKS

- What results were seen in clinical studies of WINREVAIR? What could these results mean for me?
- Were there people like me included in the clinical studies for WINREVAIR?
- Could adding WINREVAIR to my current medication plan help me reach the treatment goals that are important to me?
- What are the possible side effects of WINREVAIR? What side effects would be of concern for me?
- How would you weigh the potential benefits vs risks of WINREVAIR for someone like me?

Notes:

IMPORTANT SAFETY INFORMATION

Who should not take WINREVAIR?

- Do not take WINREVAIR if you have had a serious allergic reaction to sotatercept-csrk or any of the other ingredients in WINREVAIR.

Before taking WINREVAIR and during treatment, tell your healthcare provider:

- About all of your medical conditions and all of the medicines you take, including prescriptions and over-the-counter medicines, vitamins, or herbal supplements.
- If you are pregnant, planning to get pregnant, become pregnant, or think you may be pregnant. If you are able to get pregnant, your healthcare provider should do a pregnancy test before you start taking WINREVAIR.
- If you are breastfeeding or planning to breastfeed.

If you are able to get pregnant, you should use effective birth control during treatment with WINREVAIR and for at least 4 months after your final dose. WINREVAIR may harm your unborn baby. Do not breastfeed during treatment and for 4 months after your final dose.

Important Safety Information continued on the next page.

GETTING STARTED ON *WINREVAIR*

- What steps are involved in getting started on WINREVAIR? What would that process look like for me?
- How would I take WINREVAIR? How often would I take it?
- How would you track my results on WINREVAIR?
- How often would I come in for appointments while on WINREVAIR?

Notes:

START THE CONVERSATION WITH YOUR DOCTOR ABOUT *WINREVAIR* TODAY.

IMPORTANT SAFETY INFORMATION *(CONTINUED)*

WINREVAIR may cause serious side effects including:

- **Allergic (hypersensitivity) reactions.** Allergic reactions, including anaphylaxis, can happen within hours or days after you use WINREVAIR. Call your healthcare provider or get medical help right away if you have any of these symptoms: hives, swelling of your face, lips, tongue or throat, sudden trouble breathing or swallowing. Before your next dose of WINREVAIR, tell your healthcare provider if you have had any symptoms of an allergic reaction.
- **High level of hemoglobin in your blood.** Hemoglobin is a protein in red blood cells that carries oxygen. High levels of hemoglobin are common with WINREVAIR and can be severe and increase your risk for blood clots.
- **Severely low number of platelets in your blood.** Platelets are blood cells that help blood clot. Low platelet counts are common with WINREVAIR and can be severe and increase your risk of bleeding. Tell your healthcare provider if you develop easy bruising or bleeding, bleeding that does not stop, or nosebleeds.
- **Serious bleeding.** Serious bleeding can happen with WINREVAIR. Tell your healthcare provider if you develop any signs or symptoms of bleeding, including: vomiting blood or your vomit looks like coffee-grounds, pink or brown urine, red or black stools that look like tar, coughing up blood or blood clots, persistent headaches, nausea, dizziness or feeling weak, persistent abdominal cramps, severe back pain, or abnormally heavy menstrual bleeding.
- **Decreased fertility:** WINREVAIR may affect female and male fertility, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

Your healthcare provider will do a **blood test** before your first 5 doses of WINREVAIR, longer if needed, and then from time to time to check your levels of hemoglobin and platelets.

The most common side effects of WINREVAIR include infections, headache, nosebleeds, rash, tiny blood vessels that look like pink or red lines on the skin (spider veins), diarrhea, dizziness, redness, and bleeding gums.

These are not all the possible side effects of WINREVAIR. Call your doctor for medical advice about side effects.

Taking WINREVAIR:

- If your healthcare provider decides that you or a caregiver can give WINREVAIR at home, do not try to inject WINREVAIR until you or a caregiver have received training from your healthcare provider on the right way to prepare and inject the medicine.
- Your healthcare provider may delay treatment or change your dose if needed, based on the results of your blood tests to check your hemoglobin and platelet levels.
- See the detailed "Instructions for Use" booklet that comes with WINREVAIR for information on how to prepare and inject a dose of WINREVAIR.

Please read the accompanying Patient Information and Instructions for Use (1-vial kit, 2-vial kit) for WINREVAIR and discuss them with your doctor. The physician Prescribing Information also is available.

MERCK

Copyright © 2026 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.
Intended only for residents of the United States and its territories, including Puerto Rico.
US-SOT-03051 10/26

WINREVAIR™
(sotatercept-csrk) for injection 45 mg, 60 mg